

ORIGINAL ARTICLE

# Impact of Slow Stroke Back Massage on Blood Pressure Among Individuals With Primary Hypertension: A Quasi-Experimental Study.

Wrestina Ari Bastian <sup>1</sup>, Setiyo Adi Nugroho <sup>2</sup>, Damon Wicaksi <sup>3\*</sup>

<sup>1</sup> Universitas Nurul Jadid, Indonesia

<sup>2</sup> Universitas Nurul Jadid, Indonesia

<sup>3</sup> Universitas Bondowoso, Indonesia

\* Corresponding Author: damonwicaksi75@gmail.com

## ARTICLE INFORMATION

### Article history

Received: 2025-05-08

Revised : 2025-06-18

Accepted: 2025-06-23

### Keywords

Slow Stroke Back Massage;  
 Blood Pressure; Hypertension;  
 Non-Pharmacological  
 Intervention; Nursing Therapy

## ABSTRACT

**Introduction:** Hypertension is a major global health issue contributing significantly to cardiovascular morbidity and mortality. In Indonesia, the increasing prevalence of hypertension highlights the need for effective, low-cost, non-pharmacological interventions, such as Slow Stroke Back Massage (SSBM), which may offer physiological and psychological benefits. **Objectives:** This study aimed to determine the effect of Slow Stroke Back Massage on reducing systolic and diastolic blood pressure among individuals with primary hypertension. **Methods:** A quasi-experimental study using a non-equivalent control group pretest-posttest design was conducted. Participants aged 40–70 years with primary hypertension were purposively selected and divided into intervention and control groups. The intervention group received SSBM once daily for three consecutive days, while the control group received standard care. Blood pressure measurements were recorded before and after intervention using calibrated digital sphygmomanometers. Data were analyzed using paired and independent t-tests. **Results:** The intervention group experienced a significant reduction in both systolic (from  $150.2 \pm 10.5$  mmHg to  $137.6 \pm 9.8$  mmHg) and diastolic (from  $95.8 \pm 6.2$  mmHg to  $87.3 \pm 5.6$  mmHg) blood pressure ( $p < 0.001$ ). No statistically significant changes were observed in the control group. These results support the efficacy of SSBM as an adjunctive therapy in hypertension management. **Conclusions:** SSBM is an effective, low-cost, and non-invasive complementary intervention for lowering blood pressure in patients with primary hypertension. It can be integrated into nursing care practices, especially in primary healthcare settings, to enhance hypertension management outcomes.

Adult Health Nursing Journal is a peer-reviewed journal published by Fakultas Kesehatan, Universitas Nurul Jadid, Probolinggo, East Java.

Website: <https://fkes.unuja.ac.id>

E-mail: [ahnj@unuja.ac.id](mailto:ahnj@unuja.ac.id)

DOI : <https://doi.org/10.33650/ahnj.v2i1.11277>

This is an open access article under the [CC BY-NC-SA 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/) license



## A. Introduction

In recent decades, the global health landscape has shifted significantly, with non-communicable diseases, particularly hypertension, becoming increasingly prevalent among populations of all socioeconomic strata (Rahajeng & Tuminah, 2009; WHO, 2023). Once considered a disease of affluence, hypertension now poses a serious public health threat to low- and middle-income communities due to widespread adoption of unhealthy lifestyles (Tedjasukmana, 2012; Matyas et al., 2011). These lifestyles include poor dietary habits, physical inactivity, tobacco use, and excessive alcohol consumption—all of which contribute

to elevated cardiovascular risks (ESH, 2013; Smeltzer & Bare, 2002). Consequently, hypertension has emerged as one of the most significant contributors to global mortality and morbidity (WHO, 2023).

According to the World Health Organization (WHO), it is projected that by 2025, 1.56 billion adults worldwide will suffer from hypertension, with a disproportionate burden in low- and middle-income countries (WHO, 2023). These regions face structural challenges in hypertension management, including limited access to standardized treatment protocols, undertrained healthcare personnel, and inadequate medication supplies (Prihandana, 2012). As a result, approximately 75% of hypertensive patients fail to achieve adequate blood pressure control, leading to complications such as stroke and ischemic heart disease (Matyas et al., 2011; Moeini et al., 2011). In Southeast Asia alone, hypertension contributes to an estimated 1.5 million deaths annually (WHO, 2023).

Indonesia, as one of the most populous countries in Southeast Asia, faces an alarming rate of hypertension with prevalence exceeding national averages in several provinces, including East Java (Kemenkes RI, 2013; Depkes RI, 2007). Studies report that lifestyle factors, particularly dietary sodium intake and stress, play a critical role in the persistence of high blood pressure across the Indonesian population (Martiningsih, 2011; Smeltzer & Bare, 2002). Despite the availability of pharmacologic therapies, treatment adherence remains suboptimal due to side effects, cost, and patient perceptions (Prihandana, 2012; Brashers, 2006). This underscores the urgent need for complementary approaches that are culturally appropriate and accessible.

Among various non-pharmacologic interventions, massage therapy—particularly Slow Stroke Back Massage (SSBM)—has garnered attention as a feasible and cost-effective strategy to reduce blood pressure (Givi, 2013; Moeini et al., 2011). SSBM involves slow, rhythmic strokes across the back, which are thought to stimulate the parasympathetic nervous system, leading to muscle relaxation and physiological downregulation (Meek, 2014; Lindquist, 2005). Previous studies have shown that massage therapy can significantly reduce both systolic and diastolic blood pressure in prehypertensive and hypertensive individuals (Adib-Hajbaghery & Adib, 2014; Holland & Pokorny, 2001). These findings suggest that SSBM may serve as an effective adjunct to conventional hypertension treatment.

The physiological mechanisms underlying the benefits of SSBM are primarily attributed to its effects on the autonomic nervous system and hormonal regulation (Sylvia & Wilson, 2006; Aaronson, 2010). By stimulating mechanoreceptors in the skin, massage induces the release of endorphins and reduces cortisol levels, thereby mitigating the effects of chronic stress on cardiovascular health (Muttaqin, 2012; Moeini et al., 2011). Additionally, SSBM may promote peripheral vasodilation and improve blood circulation, resulting in decreased vascular resistance (Poetter & Perry, 2006; Smeltzer & Bare, 2002). These physiological responses align with current models of stress reduction and cardiovascular regulation.

Despite growing evidence supporting the efficacy of massage therapy, its integration into standard hypertension care remains limited, especially in primary healthcare settings (Givi, 2013; Olney, 2015). Barriers include lack of awareness among clinicians, insufficient training, and the absence of standardized protocols for implementation (Meek, 2014). Therefore, more rigorous, context-specific research is needed to establish the therapeutic validity of SSBM and to develop implementation strategies that are both effective and scalable (Holland & Pokorny, 2001; Lindquist, 2005). Furthermore, evaluating patient outcomes related to SSBM can enhance the credibility of holistic interventions in clinical practice.

As hypertension is often asymptomatic, early and sustained interventions are critical to preventing long-term complications such as myocardial infarction, stroke, and renal failure (Price et al., 2006; Kemenkes RI, 2013). Evidence suggests that combining lifestyle modification with supportive therapies such as SSBM may optimize patient adherence and

health outcomes (Ghasempour et al., 2011; Sylvia & Wilson, 2006). Moreover, the simplicity and non-invasive nature of massage therapy make it particularly suitable for community-level health promotion (Poetter & Perry, 2006; Olney, 2015). This is especially relevant in resource-limited settings where access to healthcare services and pharmacological treatments is constrained.

Given the increasing burden of hypertension and the need for sustainable interventions, there is a compelling rationale to explore the role of SSBM in blood pressure regulation among hypertensive individuals (Meek, 2014; Moeini et al., 2011). Previous research provides a strong theoretical and empirical basis for the use of massage therapy in clinical settings, yet further investigation is necessary to determine its efficacy across diverse populations (Givi, 2013; Adib-Hajbaghery & Adib, 2014). Understanding the impact of SSBM on physiological parameters such as blood pressure could inform holistic approaches to chronic disease management (Lindquist, 2005; Aaronson, 2010). Therefore, there is a need to assess the effectiveness of SSBM as a complementary intervention for hypertension.

## B. Methods

This study applied a quasi-experimental design using a non-equivalent control group pretest-posttest approach to assess the effectiveness of Slow Stroke Back Massage (SSBM) on blood pressure in individuals with primary hypertension. The intervention group received SSBM in addition to standard care, while the control group received standard care alone. Data collection was conducted over a three-day period, with pre- and post-intervention blood pressure measurements taken daily to evaluate the short-term physiological effects of the intervention. Ethical approval was granted by the local health authority, and all participants provided written informed consent prior to enrollment.

Participants were selected using purposive sampling based on inclusion and exclusion criteria. The inclusion criteria included adults aged 40 to 70 years diagnosed with primary hypertension, who were capable of communication, not experiencing hypertensive emergencies, and willing to participate. Exclusion criteria included secondary hypertension, the presence of open wounds or skin conditions on the back, spinal deformities, use of complementary therapies, or contraindications to massage such as recent fractures. The total sample consisted of hypertensive individuals who met the criteria and were divided into intervention and control groups without random allocation.

The SSBM intervention consisted of gentle, rhythmic strokes applied to the patient's back using both hands in a structured manner, beginning at the cervical vertebrae and ending at the sacral region. The massage was performed for approximately 3 to 5 minutes per session, once daily for three consecutive days, following the technique described by Meek (2014) and adapted to clinical nursing practice. Massage was conducted by trained nurses using warm hands and unscented lotion to minimize skin friction. To maintain consistency, all procedures were performed in a quiet room with patients in a prone position, and the massage pace was maintained at 12–15 strokes per minute.

Blood pressure measurements were taken before and after each intervention using a standardized, calibrated digital sphygmomanometer. Measurements included both systolic and diastolic pressures, and were recorded with the patient in a seated and rested position, following at least five minutes of inactivity. To minimize measurement bias, blood pressure was measured at the same time each day and by the same operator for each subject. Data from the control group were collected in parallel using the same protocol, except that no massage intervention was applied.

Data were analyzed using SPSS software, and statistical tests were applied to determine the effectiveness of SSBM. Descriptive statistics were used to summarize demographic and clinical characteristics, while paired t-tests were used to compare pre- and post-intervention values within groups. Independent t-tests were applied to assess differences between the

intervention and control groups. A p-value of less than 0.05 was considered statistically significant, and effect size was calculated to assess the magnitude of treatment effect.

### C. Results and Discussion

Table 1 presents the comparison of systolic and diastolic blood pressure (SBP and DBP) between the intervention group (receiving Slow Stroke Back Massage) and the control group. The intervention group showed a significant reduction in both SBP and DBP after three days of therapy. In contrast, the control group did not experience statistically significant changes. The p-values indicate that the observed changes in the intervention group were statistically significant ( $p < 0.001$ ), while those in the control group were not.

**Table 1. Comparison of Blood Pressure Pre- and Post-Intervention**

Group	Pre-test SBP (Mean $\pm$ SD)	Post-test SBP (Mean $\pm$ SD)	Pre-test DBP (Mean $\pm$ SD)	Post-test DBP (Mean $\pm$ SD)	p-value (SBP)	p-value (DBP)
Intervention	150.2 $\pm$ 10.5	137.6 $\pm$ 9.8	95.8 $\pm$ 6.2	87.3 $\pm$ 5.6	< 0.001	< 0.001
Control	148.9 $\pm$ 11.2	147.1 $\pm$ 10.7	94.7 $\pm$ 7.1	93.9 $\pm$ 6.8	0.082	0.095

These findings suggest that Slow Stroke Back Massage is effective in significantly reducing both systolic and diastolic blood pressure among patients with primary hypertension. No clinically meaningful changes were observed in the control group.

The present study demonstrates that Slow Stroke Back Massage (SSBM) significantly reduces both systolic and diastolic blood pressure in individuals with primary hypertension. This supports previous findings by Moeini et al. (2011) who reported that massage therapy produced measurable decreases in blood pressure among prehypertensive women. Physiologically, massage stimulates parasympathetic activity which promotes relaxation and vasodilation (Meek, 2014). These mechanisms contribute to reductions in vascular resistance and cardiac workload.

The magnitude of the blood pressure decrease observed in the intervention group aligns with findings by Givi (2013), who documented similar improvements in hypertensive patients receiving massage. SSBM activates pressure receptors in the skin that influence baroreceptor reflexes, leading to lowered sympathetic tone (Sylvia & Wilson, 2006). This effect enhances homeostasis of the cardiovascular system and may have long-term implications for hypertension management. In contrast, the control group in this study did not show statistically significant improvement.

From a theoretical standpoint, SSBM aligns with the Roy Adaptation Model which posits that positive sensory input can improve physiological regulation (Roy, 2009). Through physical touch and rhythm, massage delivers both physiological and psychological stimuli that can lower stress-induced blood pressure (Poetter & Perry, 2006). This supports the view that nursing interventions grounded in holistic principles can be as effective as pharmacological ones in some contexts. Such interventions can especially benefit resource-limited settings where access to medications is restricted.

The statistical significance of our findings ( $p < 0.001$ ) emphasizes the robustness of the intervention. A reduction of more than 10 mmHg in systolic blood pressure can substantially lower the risk of cardiovascular disease by up to 30% (Chobanian et al., 2003). This underscores the clinical relevance of integrating massage therapy into hypertension management protocols. Moreover, the low cost and ease of implementation make it an attractive option for public health strategies.

Notably, this study also contributes to the growing literature advocating for non-pharmacologic treatments in chronic disease management. The European Society of

Hypertension (ESH) recommends lifestyle modification as the first-line strategy for blood pressure control (ESH, 2013). However, implementation of such interventions is often inconsistent due to lack of trained personnel and structured protocols. This study demonstrates a practical, nurse-led approach that can be standardized and taught in clinical settings.

Psychological factors also play a key role in hypertension, and SSBM may exert beneficial effects by reducing anxiety and improving emotional well-being. According to Adib-Hajbaghery, Abasi, and Rajabi-Beheshtabad (2014), whole-body massage significantly reduced anxiety levels and stabilized vital signs among cardiac patients. This neuroendocrine modulation supports the blood pressure-lowering effects of massage through reductions in stress-related sympathetic activity. Therefore, massage therapy may be viewed not only as a mechanical intervention but also as a psychophysiological therapy.

In addition, the simplicity of SSBM allows for patient and caregiver education, which can empower individuals to manage their own blood pressure at home. Teaching family members to perform basic massage techniques may extend the benefits beyond the clinical environment. This could enhance continuity of care and reduce the burden on healthcare providers. However, the need for proper training and supervision remains essential to ensure safety and efficacy.

While the findings are promising, the study is not without limitations. The quasi-experimental design and non-random allocation may introduce selection bias. Future studies using randomized controlled trials (RCTs) are recommended to validate these results. Moreover, long-term follow-up would be necessary to assess the sustainability of blood pressure reduction.

In conclusion, SSBM is an effective, low-cost, and non-invasive intervention for reducing blood pressure in hypertensive individuals. It can be integrated into holistic nursing care models and utilized in primary healthcare settings to improve cardiovascular outcomes. Further research should explore its long-term effects and potential for community-based application. The findings affirm the value of evidence-based complementary therapies in modern nursing practice.

#### **D. Conclusion**

This study demonstrated that Slow Stroke Back Massage (SSBM) is an effective non-pharmacological intervention for reducing both systolic and diastolic blood pressure in individuals with primary hypertension. The intervention produced statistically and clinically significant improvements over a short period, reinforcing the physiological benefits of tactile stimulation and parasympathetic activation. Given its simplicity, low cost, and minimal risk, SSBM can serve as a valuable complementary strategy alongside standard hypertension management.

The findings support previous literature emphasizing the role of relaxation-based therapies in cardiovascular regulation and highlight the importance of holistic nursing approaches in chronic disease care. As hypertension remains a leading cause of morbidity and mortality globally, integrating massage therapy into primary care services may enhance treatment outcomes and promote patient-centered care. Further research with randomized designs and longer follow-up is recommended to confirm these effects and to assess the long-term sustainability of blood pressure reduction through massage interventions.

#### **E. Recommendations**

Based on the findings of this study, it is recommended that healthcare professionals—particularly nurses—consider incorporating Slow Stroke Back Massage (SSBM) into the



routine care of patients with primary hypertension. Training programs should be developed to equip nurses and caregivers with the necessary skills to perform SSBM safely and effectively, especially in primary healthcare and community settings.

Policymakers and healthcare administrators are encouraged to support the integration of complementary therapies such as SSBM into national hypertension control strategies. This approach can enhance patient engagement, reduce reliance on pharmacological treatment alone, and improve the overall quality of care. Additionally, public health initiatives should promote awareness of non-pharmacologic interventions for hypertension management.

Future research should focus on large-scale randomized controlled trials with extended follow-up periods to evaluate the long-term effects and cost-effectiveness of SSBM. Studies exploring patient perspectives, adherence, and the physiological mechanisms underlying the intervention will provide deeper insights and help refine clinical guidelines. Collaboration between nursing, medical, and public health disciplines is essential to advance the application of holistic care in chronic disease management.

## References

- Adib-Hajbaghery, M., Abasi, A., & Rajabi-Beheshtabad, R. (2014). Whole body massage for reducing anxiety and stabilizing vital signs of patients in cardiac care unit. *Medical Journal of the Islamic Republic of Iran*, 28, 47. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4061595/>
- Aaronson, P. I., Ward, J. P. T., & Connolly, M. J. (2010). *The cardiovascular system at a glance* (3rd ed.). Wiley-Blackwell.
- Brashers, V. L. (2006). *Clinical application of pathophysiology: An evidence-based approach*. Mosby Elsevier.
- Chobanian, A. V., Bakris, G. L., Black, H. R., Cushman, W. C., Green, L. A., Izzo, J. L., ... & National High Blood Pressure Education Program Coordinating Committee. (2003). The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *JAMA*, 289(19), 2560–2572. <https://doi.org/10.1001/jama.289.19.2560>
- Depkes RI. (2007). *Laporan Riset Kesehatan Dasar Provinsi Jawa Timur 2007*. Jakarta: Departemen Kesehatan Republik Indonesia.
- European Society of Hypertension. (2013). 2013 ESH/ESC guidelines for the management of arterial hypertension. *Journal of Hypertension*, 31(7), 1281–1357. <https://doi.org/10.1097/01.hjh.0000431740.32696.cc>
- Ghasempour, A., Moeini, M., Givi, M., & Sadeghi, T. (2011). The effect of massage therapy on blood pressure of women with prehypertension. *Iranian Journal of Nursing and Midwifery Research*, 16(1), 61–70.
- Givi, M. (2013). Durability of the effect of massage therapy on blood pressure. *International Journal of Preventive Medicine*, 4(5), 511–516.
- Holland, B., & Pokorny, M. E. (2001). Slow stroke back massage: Its effect on patients in a rehabilitation setting. *Rehabilitation Nursing*, 26(5), 182–186. <https://doi.org/10.1002/j.2048-7940.2001.tb02037.x>
- Kementerian Kesehatan Republik Indonesia. (2013). *Riset Kesehatan Dasar (Riskesdas) 2013*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan.

- Lindquist, R., Snyder, M., & Tracy, M. F. (2005). *Complementary and alternative therapies in nursing* (5th ed.). Springer Publishing Company.
- Matyas, E., Kis, O., Gombos, K., & Nagy, G. (2011). Benefit assessment of salt reduction in patients with hypertension. *Journal of Hypertension*, 29(9), 821–828.
- Meek, S. S. (2014). Effect of slow stroke back massage on relaxation in hospice clients. *Journal of Hospice and Palliative Nursing*, 16(1), 17–21. <https://doi.org/10.1097/NJH.0000000000000032>
- Moeini, M., Ghasempour, A., Givi, M., & Sadeghi, T. (2011). The effect of massage therapy on blood pressure of women with pre-hypertension. *Iranian Journal of Nursing and Midwifery Research*, 16(1), 61–70.
- Muttaqin, A. (2012). *Asuhan keperawatan pada gangguan sistem kardiovaskuler*. Jakarta: Salemba Medika.
- Olney, C. M. (2015). *Back massage: Long-term effects and dosage determination for persons with prehypertension and hypertension* (Doctoral dissertation, University of South Florida). University of South Florida Scholar Commons. <https://scholarcommons.usf.edu/etd/5625/>
- Poetter, P. A., & Perry, A. G. (2006). *Fundamentals of nursing: Concepts, process, and practice* (4th ed.). Jakarta: EGC.
- Prihandana, R. (2012). *Studi fenomenologi pengalaman kepatuhan perawatan mandiri pada pasien hipertensi di Poliklinik RSI Siti Hajar Kota Tegal* [Master's thesis, Universitas Indonesia]. Fakultas Ilmu Keperawatan UI.
- Rahajeng, E., & Tuminah, S. (2009). Prevalensi hipertensi dan determinannya di Indonesia. *Media Penelitian dan Pengembangan Kesehatan*, 19(4), 580–587. <https://dss.kalselprov.go.id/Jurnal/artikel3.pdf>
- Roy, C. (2009). *The Roy adaptation model*. Pearson.
- Smeltzer, S. C., & Bare, B. G. (2002). *Buku ajar keperawatan medikal bedah*: Brunner & Suddarth (Vol. 2, 8th ed.). Jakarta: EGC.
- Sylvia, A. P., & Wilson, L. (2006). *Patofisiologi: Konsep klinis proses-proses penyakit*. Jakarta: EGC.
- Tedjasukmana, P. (2012). *Tata laksana hipertensi*. Jakarta: Pusat Penerbitan Ilmu Penyakit Dalam, Fakultas Kedokteran UI.
- World Health Organization (WHO). (2023). *Hypertension: Factsheet*. <https://www.who.int/news-room/fact-sheets/detail/hypertension>