

# Mothers' Knowledge of Breast Cancer and Anxiety Levels in Preoperative Patients

Dodik Hartono

Universitas Hafshawaty Zainul Hasan Probolinggo, Indonesia  
Corresponding Author: ners.dodikhartono@gmail.com

## ARTICLE INFORMATION

### Article history

Received: 2025/12/27

Revised : 2025/12/29

Accepted: 2025/12/30

### Keywords

Knowledge; Anxiety level; Breast cancer; Preoperative patients; Health education

### How to cite

Hartono, D. (2025). Mothers' Knowledge of Breast Cancer and Anxiety Levels in Preoperative Patients. *Adult Health Nursing Journal*, 2(2), 32–38.  
<https://doi.org/10.33650/ahnj.v2i2.13890>

## ABSTRACT

**Background:** Breast cancer is one of the most feared diseases among women due to its uncertain etiology, limited preventive measures, unpredictable disease progression, and the potential for pain and disability in advanced stages. Limited knowledge about breast cancer and surgical procedures may contribute to anxiety among patients, particularly during the preoperative period, which can negatively affect psychological readiness and clinical outcomes. **Objective:** This study aimed to examine the relationship between mothers' knowledge of breast cancer and anxiety levels among preoperative breast cancer patients. **Methods:** A correlational study with a cross-sectional design was conducted among preoperative breast cancer patients scheduled for surgery in a central surgical unit. A total of 84 respondents were recruited using accidental sampling. Data were collected using structured questionnaires to assess knowledge levels and the Hamilton Anxiety Rating Scale (HARS) to measure anxiety. Statistical analysis was performed using Spearman's rho correlation test with a significance level of 0.05. **Results:** The analysis showed a statistically significant relationship between mothers' knowledge of breast cancer and anxiety levels among preoperative breast cancer patients ( $p = 0.013$ ), indicating that lower knowledge levels were associated with higher anxiety. **Conclusion:** There is a significant association between mothers' knowledge of breast cancer and preoperative anxiety levels. Strengthening patient education through structured health counseling and standardized operating procedures may help reduce anxiety and improve psychological preparedness before surgery.

**Adult Health Nursing Journal** is a peer-reviewed journal published by Fakultas Kesehatan, Universitas Nurul Jadid, Probolinggo, East Java.

Website: <https://fkes.unuja.ac.id>

E-mail: [adulhealthnurse@gmail.com](mailto:adulhealthnurse@gmail.com)

DOI : <https://doi.org/10.33650/ahnj.v2i2.13890>

This is an open access article under the CC BY-NC-SA 4.0 license



## A. Introduction

Breast cancer is the most common cancer among women worldwide and remains a major public health concern due to its high incidence, complex etiology, and substantial physical and psychological burden (Freihat et al., 2025; Sung et al., 2021; WHO, 2023). Despite advances in early detection and treatment, many patients continue to experience significant distress associated with diagnosis and treatment, particularly in low- and middle-income settings. The disease trajectory is often unpredictable, and advanced stages may involve severe pain, functional impairment, and reduced quality of life (Faller et al., 2013; Mystakidou et al., 2005).

Beyond its physical consequences, breast cancer has profound psychological implications. Anxiety and broader psychological distress are among the most prevalent responses following

diagnosis and during treatment, affecting patients' emotional stability and coping capacity (Mehnert et al., 2018; Tao et al., 2024). Recent systematic reviews indicate that approximately half of breast cancer patients experience clinically relevant psychological distress at some point during the disease trajectory (Tang et al., 2024; Tao et al., 2024). Psychological distress in cancer patients has been associated with poorer treatment adherence, impaired functioning, and less favorable clinical outcomes, underscoring the importance of integrating mental health into oncologic care (Hessami et al., 2025; Kim et al., 2020; Roche et al., 2023).

Surgical intervention remains a cornerstone of breast cancer management, particularly for patients with operable disease. However, the preoperative period is often characterized by heightened anxiety related to uncertainty about surgical outcomes, anesthesia, postoperative pain, and potential changes in body image (Kim et al., 2020; Tao et al., 2024). Preoperative anxiety is not merely an emotional reaction but a clinically relevant condition that can influence physiological responses and postoperative recovery. Evidence suggests that elevated preoperative anxiety is associated with increased postoperative pain, delayed recovery, and reduced patient satisfaction (Kim et al., 2020; Mulugeta et al., 2018; Tao et al., 2024).

Patients experiencing high anxiety levels may also demonstrate reduced cooperation with perioperative care and longer hospitalization. Consequently, identifying modifiable factors that contribute to preoperative anxiety is critical for improving surgical outcomes (Katsohiraki et al., 2020; Mulugeta et al., 2018; Tang et al., 2024). One important factor influencing anxiety is patients' knowledge regarding breast cancer and planned surgical procedures. Adequate knowledge enables patients to understand their condition, anticipate the treatment process, and develop effective coping strategies (Darville-Beneby et al., 2023; Vodermaier et al., 2009). Conversely, insufficient or inaccurate information may increase uncertainty, fear, and catastrophic thinking, thereby exacerbating anxiety (Sbaity et al., 2025; Zhong et al., 2021).

Several studies have demonstrated that structured preoperative education and information provision can significantly reduce anxiety and improve psychological readiness for surgery. Preconsultation educational group interventions and multimedia education have been shown to increase knowledge, reduce decisional conflict, and lower state anxiety before breast surgery or reconstruction (Sadeghi et al., 2025; Spiegel & Giese-Davis, 2003; Zhong et al., 2021). Psycho-oncological interventions that enhance patients' understanding and sense of control have also been shown to reduce emotional distress and improve quality of life among cancer patients (Faller et al., 2013; Mehnert et al., 2014). These findings highlight the importance of knowledge and information as key determinants of psychological adaptation in cancer care.

However, despite growing evidence supporting the role of patient education, empirical data examining the direct relationship between patients' knowledge levels and preoperative anxiety specifically in breast cancer surgery remain limited. Existing studies often focus on general psychological distress or decision-related outcomes (such as decisional conflict or regret) rather than knowledge-specific factors influencing anxiety (Mehnert et al., 2014; Zhong et al., 2021). Recent work has emphasized that breast cancer-related psychological distress arises from multiple interrelated risk factors, including disease-related variables, sociodemographic characteristics, and coping style, but knowledge as a distinct contributor is less well explored (Tang et al., 2024; Tao et al., 2024).

Therefore, this study aims to examine the relationship between mothers' knowledge of breast cancer and anxiety levels among patients undergoing preoperative breast cancer surgery. Understanding this relationship is expected to provide evidence-based support for developing structured educational interventions to reduce anxiety and improve psychological preparedness in surgical breast cancer care.

## B. Methods

This study employed a correlational design with a cross-sectional approach to examine the relationship between mothers' knowledge of breast cancer and anxiety levels among preoperative breast cancer patients. The cross-sectional design allowed for the simultaneous assessment of knowledge and anxiety at a single point in time, making it appropriate for identifying associations between variables without intervention or follow-up.

The study population consisted of breast cancer patients scheduled for surgical treatment in a central surgical unit. A total of 84 respondents were included in the study using accidental sampling, whereby eligible patients who met the inclusion criteria and were available during the data collection period were recruited. Inclusion criteria comprised adult female patients diagnosed with breast cancer, scheduled for elective surgery, and able to communicate effectively. Patients with severe cognitive impairment or diagnosed psychiatric disorders were excluded.

Two main variables were assessed in this study: knowledge of breast cancer as the independent variable and anxiety level as the dependent variable. Knowledge was defined as patients' understanding of breast cancer, including its nature, treatment options, and surgical procedures. Anxiety was defined as an emotional state characterized by feelings of tension, worry, and physiological arousal experienced prior to surgery.

Data were collected using validated instruments. Patients' knowledge of breast cancer was measured using a structured questionnaire developed based on relevant literature and clinical guidelines. Anxiety levels were assessed using the Hamilton Anxiety Rating Scale (HARS), a widely used and validated instrument for measuring anxiety severity in clinical populations. Higher scores on the HARS indicate greater levels of anxiety.

Prior to data collection, ethical approval was obtained from the relevant institutional authority. All participants received an explanation of the study objectives and procedures and provided informed consent. Confidentiality and anonymity were strictly maintained, and participants were informed of their right to withdraw from the study at any time without consequences.

Data analysis was performed using the Statistical Package for the Social Sciences (SPSS). Descriptive statistics were used to summarize respondents' characteristics and variable distributions. The relationship between knowledge levels and anxiety scores was analyzed using Spearman's rho correlation test, as the data were not assumed to be normally distributed. Statistical significance was set at a p-value of less than 0.05.

## C. Results

A total of 84 preoperative breast cancer patients participated in this study. Most respondents were aged over 40 years (61.9%), had a senior high school educational background (58.3%), and were housewives (52.4%). These characteristics indicate that the majority of participants were middle-aged women with moderate educational attainment.

Regarding knowledge of breast cancer, 45.2% of respondents demonstrated a moderate level of knowledge, while 32.1% had good knowledge and 22.7% had poor knowledge. Anxiety levels assessed using the Hamilton Anxiety Rating Scale (HARS) showed that 48.8% of respondents experienced mild anxiety, 38.1% moderate anxiety, and 13.1% severe anxiety.

Cross-tabulation analysis demonstrated that patients with good knowledge predominantly experienced mild anxiety, whereas those with poor knowledge were more likely to experience moderate to severe anxiety.

**Table 1: Distribution of Knowledge Levels and Anxiety Among Preoperative Breast Cancer Patients**

Variable	Category	n	%
Knowledge	Good	27	32.1
	Moderate	38	45.2
	Poor	19	22.7
Anxiety	Mild	41	48.8
	Moderate	32	38.1
	Severe	11	13.1

Spearman's rho correlation analysis revealed a statistically significant positive correlation between knowledge level and anxiety severity ( $p = 0.013$ ), indicating that lower knowledge levels were associated with higher anxiety among preoperative breast cancer patients.

**Table 2: Association Between Knowledge Level and Anxiety Severity**

Knowledge Level	Mild Anxiety	Moderate Anxiety	Severe Anxiety	Total
Good	21 (25.0%)	6 (7.1%)	0 (0.0%)	27 (32.1%)
Moderate	18 (21.4%)	17 (20.2%)	3 (3.6%)	38 (45.2%)
Poor	2 (2.4%)	9 (10.7%)	8 (9.6%)	19 (22.7%)
<b>Total</b>	<b>41 (48.8%)</b>	<b>32 (38.1%)</b>	<b>11 (13.1%)</b>	<b>84 (100%)</b>

#### D. Discussion

The results indicate that nearly half of the respondents had a moderate level of knowledge regarding breast cancer, while more than one-fifth demonstrated poor knowledge. This finding suggests that despite increasing public awareness of breast cancer, substantial gaps in patient understanding remain, which is consistent with recent reports that many patients still experience limited health literacy regarding cancer and its treatments (Mehnert et al., 2014; Zhong et al., 2021). Limited health literacy may hinder patients' ability to interpret medical information accurately and to engage effectively in shared decision-making.

Anxiety was prevalent among preoperative breast cancer patients, with more than half experiencing moderate to severe anxiety. This finding aligns with previous studies reporting that anticipation of surgery, fear of anesthesia, uncertainty about prognosis, and concerns about body image contribute significantly to preoperative anxiety (Faller et al., 2013; Mehnert et al., 2014). Elevated anxiety in the perioperative period has been shown to adversely affect pain perception, recovery, and satisfaction with care (Mulugeta et al., 2018).

The significant correlation between knowledge level and anxiety supports the hypothesis that insufficient understanding of breast cancer is associated with higher anxiety. Patients with good knowledge predominantly experienced mild anxiety, whereas those with poor knowledge were more likely to report severe anxiety. This pattern is consistent with evidence that inadequate or unclear information exacerbates psychological distress in cancer patients, whereas appropriate information and counseling can mitigate anxiety and improve emotional adjustment (Faller et al., 2013; Mehnert et al., 2014).

From a psychological perspective, knowledge enhances patients' perceived control and coping capacity. Patients who understand their disease and treatment options are better able to anticipate surgical procedures, set realistic expectations, and manage emotional responses. Conversely, lack of knowledge may promote uncertainty and catastrophic thinking, thereby intensifying anxiety (Mehnert et al., 2018; Tao et al., 2024). This is in line with cognitive theories of anxiety, which emphasize the role of threat appraisal and perceived controllability in the development of anxious responses.

These findings highlight the importance of structured preoperative education as part of comprehensive breast cancer care. Psycho-oncological and educational interventions have been shown to reduce emotional distress, improve psychological preparedness for surgery,

and enhance overall quality of life in patients with cancer (Faller et al., 2013; Mehnert et al., 2014). Nurses and healthcare providers play a central role in delivering accurate, empathetic, and individualized information tailored to patients' educational levels and informational needs (Sadeghi et al., 2025).

The larger sample size in this study strengthens the reliability of the findings compared with smaller studies, increasing the precision of estimated associations. However, the cross-sectional design limits causal inference, and anxiety levels were measured at a single time point before surgery. Longitudinal studies are recommended to assess changes in anxiety across the perioperative period and to evaluate the long-term impact of educational interventions on psychological outcomes (Mehnert et al., 2018; Tao et al., 2024).

Overall, this study confirms that patients' knowledge of breast cancer is significantly associated with anxiety levels during the preoperative phase. Enhancing patient education through standardized, evidence-based health education protocols may reduce anxiety and improve psychological readiness for breast cancer surgery, and should be considered an integral component of patient-centered breast cancer care (Kim et al., 2020; Mulugeta et al., 2018; Tao et al., 2024).

### **Implication and limitation**

The findings of this study indicate that patients' knowledge of breast cancer plays a significant role in influencing preoperative anxiety levels, where lower knowledge is associated with higher anxiety. This highlights the importance of integrating structured patient education into preoperative care to improve psychological preparedness and reduce emotional distress. Healthcare providers, particularly nurses, should prioritize delivering clear, accurate, and patient-centered information regarding disease conditions and surgical procedures. However, this study has several limitations. The cross-sectional design limits the ability to establish causal relationships between knowledge and anxiety. The use of accidental sampling may introduce selection bias, and the study was conducted in a single setting, which may limit generalizability. Additionally, anxiety was measured at a single time point, which may not fully capture fluctuations during the perioperative period. Future studies using longitudinal designs and broader populations are recommended to strengthen the evidence.

### **Relevance for Practice**

This study provides important practical implications for improving preoperative care in breast cancer patients by emphasizing the role of patient education in reducing anxiety. Healthcare professionals should implement structured educational interventions, including preoperative counseling and standardized information delivery, to enhance patients' understanding and coping abilities. By improving knowledge, nurses can help reduce anxiety levels, increase patient confidence, and support better psychological readiness for surgery, ultimately contributing to improved clinical outcomes and patient satisfaction.

## **E. Conclusion**

This study demonstrates a statistically significant relationship between mothers' knowledge of breast cancer and anxiety levels among preoperative breast cancer patients. Patients with lower levels of knowledge were more likely to experience moderate to severe anxiety prior to surgery, whereas those with better knowledge predominantly reported mild anxiety. These findings indicate that knowledge plays an important role in shaping patients' psychological responses during the preoperative phase. Enhancing patients' understanding of breast cancer and surgical procedures is therefore essential to reduce anxiety and improve psychological readiness for surgery.

### Acknowledgment

The authors would like to express their sincere gratitude to all parties who supported the completion of this study. The authors also thank all participants who willingly contributed their time and responses to this research.

### Author Contribution

Dodik Hartono contributed to the study conceptualization, research design, data collection, data analysis, interpretation of results, and manuscript writing. The author has read and approved the final version of the manuscript and agrees to be accountable for all aspects of the work.

### Funding

This research received no external funding.

### Declaration of Conflicting Interest

The authors declare no conflict of interest.

### Declaration of Use of AI in Scientific Writing

The authors declare that generative AI and AI-assisted technologies were used to support language editing and grammatical refinement of the manuscript.

### References

- Darville-Beneby, R., Lomanowska, A. M., Yu, H. C., Jobin, P., Rosenbloom, B. N., Gabriel, G., Daudt, H., Negraeff, M., Di Renna, T., Hudspith, M., & Clarke, H. (2023). The Impact of Preoperative Patient Education on Postoperative Pain, Opioid Use, and Psychological Outcomes: A Narrative Review. *Canadian Journal of Pain, 7*(2), 2266751. <https://doi.org/10.1080/24740527.2023.2266751>
- Faller, H., Schuler, M., Richard, M., Heckl, U., Weis, J., & Küffner, R. (2013). Effects of psycho-oncologic interventions on emotional distress and quality of life in adult patients with cancer: systematic review and meta-analysis. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology, 31*(6), 782–793. <https://doi.org/10.1200/JCO.2011.40.8922>
- Freihat, O., Sipos, D., & Kovacs, A. (2025). Global burden and projections of breast cancer incidence and mortality to 2050: a comprehensive analysis of GLOBOCAN data. *Frontiers in Public Health, 13*, 1622954. <https://doi.org/10.3389/fpubh.2025.1622954>
- Hessami, A., Ghadirzadeh, E., Ashrafi, S., Taghavi, F., Elyasi, F., Gheibi, M., Zaboli, E., Alizadeh-Navaei, R., Akbari Gelvardi, F., Hedayatizadeh-Omran, A., & Moosazadeh, M. (2025). Depression, anxiety, and stress in breast cancer patients: prevalence, associated risk factors, and clinical correlates. *Scientific Reports, 15*(1), 31084. <https://doi.org/10.1038/s41598-025-17285-7>
- Ilbawi AM. WHO Report on Cancer: Setting priorities, investing wisely and providing care for all. In: Wild CP, Weiderpass E, Stewart BW, editors. *World Cancer Report: Cancer research for cancer prevention*. Lyon (FR): International Agency for Research on Cancer; 2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK606484/>.
- Katsohiraki, M., Pouloupoulou, S., Fyrfiris, N., Koutelekos, I., Tsiotinou, P., Adam, O., Vasilopoulou, E., & Kapritsou, M. (2020). Evaluating Preoperative Anxiety Levels in Patients Undergoing Breast Cancer Surgery. *Asia-Pacific Journal of Oncology Nursing, 7*(4), 361–364. [https://doi.org/10.4103/apjon.apjon\\_31\\_20](https://doi.org/10.4103/apjon.apjon_31_20)
- Kim, J., Cho, J., Lee, S. K., Choi, E.-K., Kim, I.-R., Lee, J. E., Kim, S. W., & Nam, S.-J. (2020). Surgical impact on anxiety of patients with breast cancer: 12-month follow-up prospective longitudinal study. *Annals of Surgical Treatment and Research, 98*(5), 215–223. <https://doi.org/10.4174/ast.2020.98.5.215>
- Mehnert, A., Brähler, E., Faller, H., Härter, M., Keller, M., Schulz, H., Wegscheider, K., Weis, J., Boehncke, A., Hund, B., Reuter, K., Richard, M., Sehner, S., Sommerfeldt, S., Szalai, C., Wittchen, H.-U., & Koch, U. (2014). Four-week prevalence of mental disorders in patients with cancer across major tumor entities. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology, 32*(31), 3540–3546. <https://doi.org/10.1200/JCO.2014.56.0086>
- Mehnert, A., Hartung, T. J., Friedrich, M., Vehling, S., Brähler, E., Härter, M., Keller, M., Schulz, H., Wegscheider, K., Weis, J., Koch, U., & Faller, H. (2018). One in two cancer patients is significantly

- distressed: Prevalence and indicators of distress. *Psycho-Oncology*, 27(1), 75–82. <https://doi.org/10.1002/pon.4464>
- Mulugeta, H., Ayana, M., Sintayehu, M., Dessie, G., & Zewdu, T. (2018). Preoperative anxiety and associated factors among adult surgical patients in Debre Markos and Felege Hiwot referral hospitals, Northwest Ethiopia. *BMC Anesthesiology*, 18(1), 155. <https://doi.org/10.1186/s12871-018-0619-0>
- Mystakidou, K., Tsilika, E., Parpa, E., Katsouda, E., Galanos, A., & Vlahos, L. (2005). Assessment of Anxiety and Depression in Advanced Cancer Patients and their Relationship with Quality of Life. *Quality of Life Research*, 14(8), 1825–1833. <https://doi.org/10.1007/s11136-005-4324-3>
- Roche, K. N., Cooper, D., Armstrong, T. S., & King, A. L. (2023). The link between psychological distress and survival in solid tumor patients: A systematic review. *Cancer Medicine*, 12(3), 3343–3364. <https://doi.org/10.1002/cam4.5200>
- Sadeghi, N., Salari, N., & Jalali, R. (2025). Effect of multimedia education on anxiety and pain in patients undergoing laparoscopic cholecystectomy: a Solomon four-group randomized controlled trial. *Scientific Reports*, 15(1), 9357. <https://doi.org/10.1038/s41598-024-77207-x>
- Sbaity, E., Diab, T., Haroun, J., Ramadan, N., Khalil, G., Chamseddine, N., Diab, R., Mansour, H., El Sayed, M., Charafeddine, M., Abbas, J., & Assi, H. I. (2025). Understanding Patient Decision-Making in Breast Cancer Surgery: Risk Perception, Communication, and Psychosocial Influences. *Medical Sciences*, 13(4). <https://doi.org/10.3390/medsci13040225>
- Spiegel, D., & Giese-Davis, J. (2003). Depression and cancer: mechanisms and disease progression. *Biological Psychiatry*, 54(3), 269–282. [https://doi.org/10.1016/s0006-3223\(03\)00566-3](https://doi.org/10.1016/s0006-3223(03)00566-3)
- Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021). Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA: A Cancer Journal for Clinicians*, 71(3), 209–249. <https://doi.org/10.3322/caac.21660>
- Tang, W.-Z., Mangantig, E., Iskandar, Y. H. P., Cheng, S.-L., Yusuf, A., & Jia, K. (2024). Prevalence and associated factors of psychological distress among patients with breast cancer: a systematic review and meta-analysis. *BMJ Open*, 14(9), e077067. <https://doi.org/10.1136/bmjopen-2023-077067>
- Tao, L., Xiang, Y., Zeng, X., Fu, L., Li, J., & Chen, H. (2024). Incidence and risk factors for psychological distress in adult female patients with breast cancer: a systematic review and meta-analysis. *Frontiers in Psychiatry*, Volume 15-2024. <https://doi.org/10.3389/fpsy.2024.1309702>
- Vodermaier, A., Linden, W., & Siu, C. (2009). Screening for emotional distress in cancer patients: a systematic review of assessment instruments. *Journal of the National Cancer Institute*, 101(21), 1464–1488. <https://doi.org/10.1093/jnci/djp336>
- Zhong, T., Quong, W. L., Cheng, T., Kerrebijn, I., Butler, K., Hofer, S. O. P., O'Neill, A. C., Cil, T. D., & Metcalfe, K. A. (2021). Preconsultation Educational Group Intervention Can Address the Knowledge Gap in Postmastectomy Breast Reconstruction. *Annals of Plastic Surgery*, 86(6), 695–700. <https://doi.org/10.1097/SAP.0000000000002603>